

PROCESS FOR THE RACEMISATION OF ENANTIOMERICALLY ENRICHED

5

 $\alpha$ -AMINO NITRILES

The invention relates to a process for the racemisation of enantiomerically enriched  $\alpha$ -amino nitriles.

Many enantiomerically enriched active ingredients like pharmaceuticals and agrochemicals can be prepared using enantiomerically enriched  $\alpha$ -amino nitriles, which are commonly used as a precursor for  $\alpha$ -amino acids or derivatives thereof. The preparation of pharmaceuticals or agrochemicals generally involves a process with many consecutive steps.  $\alpha$ -Amino nitriles can be converted into a wide variety of compounds, for example  $\alpha$ -amino acids,  $\alpha$ -amino acid amides,  $\beta$ -amino alcohols, vicinal diamines and protected  $\alpha$ -amino aldehydes, and therefore are broadly applicable in the preparation of enantiomerically enriched active ingredients. The preparation of enantiomerically enriched compounds commonly includes a resolution step wherein the undesired enantiomer is left as a byproduct. In such resolutions the theoretical maximum yield with regard to the starting material is 50%. The yield can increase significantly by racemisation and recycling of the undesired enantiomer. Preferably the resolution process is combined with racemisation in situ. Therefore, there exists a need for a racemisation process of  $\alpha$ -amino nitriles that is compatible with a large variety of resolution processes.

A process wherein an in situ racemisation of an enantiomerically enriched  $\alpha$ -amino nitrile occurs is described in US-A-4683324.

A disadvantage of the process of US-A-4683324 is that a protic solvent has to be used. It is known that  $\alpha$ -amino nitriles partly decompose in protic solvents. Therefore, in the process of US-A-4683324 HCN is added in order to suppress decomposition. From EP-A-1050529 it is further known that non polar, (but actually meaning aprotic) solvents stabilize  $\alpha$ -amino nitriles with respect to decomposition and prevent racemisation.

It is the object of the invention to provide a process for the racemisation of  $\alpha$ -amino nitriles wherein a racemisation with less decomposition is achieved, and that can be used in combination with resolutions in the preparation of enantiomerically enriched compounds in high yield and high enantiomeric excess.

This object is achieved according to the invention by using a Lewis

acid catalyst. Lewis acids by definition are able to accept an electron pair from compounds that have a free available electron pair.

Suitable Lewis acid catalysts according to the invention comprise a metal ion M, which is complexed according to the general structure

5



in which n represents an integer larger than or equal to 1, for instance is equal to 1, 2, 3, 4, 5, 6, 7, ... and p, q and r each independently represent an integer larger than or equal to 0, for instance 0, 1, 2, 3, 4, 5, 6, 7, ..., and in which n and p are chosen such that  $M_nX_p$  is neutral.

10

The metal M may be in the ionic form ( $M^{k+}$ ) with suitable counter ions X, wherein k represents the valence of the metal ion, or in the zero valent form ( $M^0$ ) with p = 0.

15

Suitable metals are for instance metals chosen from main group elements IA-IVA of the Periodic Table (CAS version), the transition metals and the lanthanides, preferably Al, Ti, Zr, or a lanthanide, for example Sm or La, or combinations thereof.

20

X represents an anionic counter ion or covalently bound anionic ligand for non zero valent metals. Suitable examples are halides, in particular  $Cl^-$  or  $Br^-$ ; alkyl groups with e.g. 1-12 C-atoms, for example methyl, ethyl, n-propyl ( $^nPr$ ) or i-butyl ( $^iBu$ ) groups, alkoxy groups with e.g. 1-12 C-atoms, for example n-pentoxy, i-propoxy, t-butoxy groups preferably the alkoxy groups derived from a secondary alcohol; anions derived from amides, aminoalcohols or amines;  $CN^-$  groups; anionic aromatic ligands, in particular cyclopentadienyl ( $Cp$ ), pentamethyl cyclopentadienyl ( $Cp^*$ ) or indenyl.

25

S represents a so-called spectator ligand, a neutral ligand which is difficult to exchange by other ligands, for example aromatic compounds or olefines. Examples of aromatic compounds are aromatic hydrocarbons, for instance benzene, toluene, cumene, cymene or mesitylene. Suitable olefines are for instance dienes, in particular 1,5-cyclooctadiene or 1,5-hexadiene.

30

L represents a neutral ligand, which can be relatively easily exchanged by other ligands, for example phosphines in particular  $PPh_3$  or  $PCy_3$ , nitriles or coordinating solvent molecules, especially tetrahydrofuran (THF), acetonitrile, dimethylformamide, alcohols, amines, in particular tertiary amines, for example  $Et_3N$ .

35

Catalyst represented by the general structure  $M_nX_pS_qL_r$  as described above are generally known in the art. As will be understood by the person skilled in the art the distinction between spectator ligand S and neutral ligands L is gradual and

amenable to discussion; besides that it may depend on the reaction conditions.

Examples of suitable catalyst are: LiBr, Al(O<sup>i</sup>Pr)<sub>3</sub>, AlCl<sub>3</sub>, Al(alkyl)<sub>3</sub>, with each alkyl is a C1-C12 alkyl group, Al(Et<sub>2</sub>)CN, Al(<sup>i</sup>Bu)<sub>2</sub>H,  $\{ \text{Al}(\text{CH}_3)\text{-O} \}_n$ , Zr(O<sup>i</sup>Bu)<sub>4</sub>, Zirconocene, Sm(O<sup>i</sup>Pr)<sub>3</sub>, Sm(Cp)<sub>3</sub>, La(Cp)<sub>3</sub>. Particularly preferred catalyst are aluminum  
5 alkoxides, aluminum alkyls, lanthanide alkoxides, for instance samarium alkoxides, and lanthanocenes, for instance lanthanide tris-cyclopentadienyl compounds.

The racemisation can be carried out in the presence of a solvent. Preferably aprotic solvents are used in the racemisation according to the invention, for instance aliphatic hydrocarbons, for example hexane, heptane; aromatic hydrocarbons,  
10 for example toluene and xylene; ethers, for example methyl t-butyl ether (MTBE); esters, for example ethyl acetate; or mixtures thereof. The concentration of the amino nitrile in the reaction mixture preferably is between 0.1 and 2 M.

The temperature for the racemisation preferably will be chosen as high as possible and such that still a sufficiently low level of decomposition is  
15 maintained under the reaction conditions. The optimal temperature can be determined by the skilled person and lies for instance between -20 and 120 °C, preferably between 30 and 100 °C.

The racemisation according to the invention can be combined with a separate resolution processes or with an in situ resolution process so that in principle  
20 the product can be obtained in 100% yield and 100% ee. In a particularly preferred embodiment of the invention, racemisation is applied in combination with an in situ resolution process.

Examples of resolution processes that can be combined with the racemisation according to the invention are for instance, crystallization induced  
25 resolutions (classical resolutions), for instance resolutions via diastereoisomeric salt formation or entrainment, enzymatic resolutions, or resolutions achieved by physical separation methods. Examples of resolution process in combination with an in situ racemisation are, for instance, crystallization induced asymmetric transformations or dynamic kinetic resolutions (DKR). A clear advantage of the racemisation according to  
30 the invention is that an aprotic solvent can be chosen. In order to achieve a "100% yield, 100° ee" result as a rule it is better to use aprotic solvents in resolution processes. For instance, aprotic organic solvents show less deactivation of enzymes in enzymatic processes than protic organic solvents. Thus, in combination with an enzymatic resolution (DKR) aprotic solvents are particularly suitable solvents to be

used in stereoselective reactions e.g. stereoselective acylation reactions or stereoselective nitrile converting reactions. For asymmetric transformations low solubilities of diastereomeric salts are preferred and can be achieved by aprotic solvents.

5 In such "100% yield and 100% ee" concepts the racemisation system is chosen such that the racemisation system is compatible with the stereo differentiating process, for instance with the resolving agents or the enzyme used in the resolution. The choice of the optimum reaction conditions will as a rule be a compromise between the optimum resolving or stereo differentiating process conditions and the optimum racemisation conditions. The person skilled in the art will be able to find the optimal conditions for his situation. The temperature range, for instance, mostly will be chosen between 0 and 120 °C, preferably 20 and 110 °C. The concentration of the amino nitrile in the reaction mixture preferably is between 0.1 and 2M.

15 Examples of amino nitriles that can be racemised according to the invention are for instance amino nitriles with formula (1)



20 wherein \* denotes a chiral C-atom, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> each independently represent H, an optionally substituted alkyl, aryl, alkaryl or aralkyl group with, for instance 1-20 C-atoms, or R<sup>1</sup> and R<sup>2</sup> or R<sup>1</sup> and R<sup>3</sup> or R<sup>3</sup> and R<sup>4</sup> together with the atom(s) to which they are attached form an optionally substituted 4-8 membered (hetero)cyclic group with for instance 3-20 C-atoms.

25 The invention will be elucidated by the following examples, without however being limited thereby.

#### Example I

##### Racemisation using diethylaluminum cyanide

30 In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 1 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. A 1 M solution of diethylaluminum cyanide

in toluene was added by syringe and the reaction mixture was stirred under nitrogen at 80°C for 18 h. The results are given in Table 1.

Table 1: Racemisation of (R)-2-amino-3-phenylpropionitrile

| Exp. | Catalyst solution (μl) | Catalyst (mol %) | E.e. (%) |
|------|------------------------|------------------|----------|
| 1    | 10                     | 10               | 0        |
| 2    | 25                     | 25               | 0        |
| 3    | 50                     | 50               | 0        |

5

### Example II

#### Racemisations at room temperature using aluminum complexes

Fresh stock solutions (0.1 m in toluene) were prepared from a) diethylaluminum cyanide ((C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>AlCN), b) triethylaluminum ((C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>Al), c) tri-  
10 butylaluminum [(CH<sub>3</sub>)<sub>2</sub>CHCH<sub>2</sub>]<sub>3</sub>Al and d) MAO (methylaluminoxane). each stock solution was used as catalyst solution in following racemisation procedure for 2-amino-3-phenylpropionitrile.

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99  
15 %) was dissolved in 0.9 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To the solution, 0.1 ml of catalyst solution was added by syringe and the reaction mixture was stirred under nitrogen at room temperature. The results are given in Table 2, below.

20 Table 2: Racemisation of (R)-2-amino-3-phenylpropionitrile

| Exp. | Catalyst   | t (h) | E.e. (%) |
|------|--|-------|----------|
| 1    | C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> AlCN                    | 1     | 0        |
| 2    | ((C <sub>2</sub> H <sub>5</sub> ) <sub>3</sub> Al                    | 1     | 0        |
| 3    | [(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> ] <sub>3</sub> Al | 0.25  | 14       |
| 4    | MAO  | 17    | 43       |

The lowest e.e. represents the best result

### Example III

#### Racemisation at room temperature using DIBAL

A 0.1 M stock solution of DIBAL (diisobutylaluminum hydride) in  
 5 hexane was prepared by dilution from a 1 M solution in hexane.  
 In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved  
 in 0.9 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture  
 was degassed by nitrogen. To the solution, 0.1 ml of the 0.1 M stock solution in hexane  
 was added by syringe and the reaction mixture was stirred under nitrogen at room  
 10 temperature. After 1.25 h, 2-amino-3-phenylpropionitrile was racemised to 59 % e.e..

### Example IV

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using heterogeneous racemisation catalyst

15 Catalyst and solid (R)-2-amino-3-phenylpropionitrile (0.1 mmol, e.e. >  
 99 %) were weight under a dry atmosphere in a glovebox in a pre-dried 5 ml vial,  
 equipped with magnetic stirrer bar. The vial was sealed with a teflon-lined cap and  
 taken out of the glovebox. To the heterogeneous mixture, 1 ml dry solvent was added  
 20 by syringe, and the reaction mixture was stirred under nitrogen at temperature given in  
 Table 3 below.

Table 3: Racemisation of (R)-2-amino-3-phenylpropionitrile using heterogeneous catalysts

| Exp. | Catalyst   | Catalyst (mg) | Solvent | T (°C) | t (h) | e.e. (%) |
|------|--|---------------|---------|--------|-------|----------|
| 1    | Sm <sub>5</sub> O[OC <sub>3</sub> H <sub>7</sub> ] <sub>13</sub> | 60            | THF     | 75     | 17    | 10       |
| 2    | Sm <sub>5</sub> O[OC <sub>3</sub> H <sub>7</sub> ] <sub>13</sub> | 1.5           | THF     | 75     | 24    | 92       |
| 3    | Sm <sub>5</sub> O[OC <sub>3</sub> H <sub>7</sub> ] <sub>13</sub> | 1.5           | Toluene | 75     | 24    | 70       |
| 4    | La(Cp) <sub>3</sub>  | 33.4          | Toluene | 80     | 1     | 0        |
| 5    | La(Cp) <sub>3</sub>  | 3.3           | Toluene | 50     | 17    | 7        |
| 6    | Sm(Cp) <sub>3</sub>  | 3.5           | Toluene | 50     | 17    | 23       |
| 7    | Al <sub>2</sub> O <sub>3</sub>                                   | 10.3          | Toluene | 60     | 24    | 35       |
| 8    | SiO <sub>2</sub>   | 24            | Toluene | 60     | 16    | 88       |

### Example V

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using aluminum isopropoxide

##### A) Catalysis with solid aluminum isopropoxide

5

Solid aluminum isopropoxide ( $\text{Al}(\text{O}^i\text{Pr})_3$ ) and solid (R)-2-amino-3-phenylpropionitrile (0.1 mmol, e.e. > 99 %) were placed in a vial of 5 ml and dissolved in 1 ml solvent. The vial was sealed with a teflon-lined cap, the solution degassed by nitrogen and the racemisation was carried out at temperatures given in Table 4 below.

10

Table 4: Racemisation of (R)-2-amino-3-phenylpropionitrile using solid  $\text{Al}(\text{O}^i\text{Pr})_3$

| Exp. | Catalyst (mg) | Solvent | T (°C) | t (h) | e.e. (%) |
|------|---------------|---------|--------|-------|----------|
| 1    | 2             | THF     | 75     | 17    | 23       |
| 2    | 10            | THF     | 75     | 17    | 0        |
| 3    | 0.8           | Toluene | 75     | 18    | 29       |
| 4    | 2             | Toluene | 20     | 70    | 83       |
| 5    | 3.4           | Toluene | 80     | 2     | 72       |

##### B) Aluminum isopropoxide prepared from other aluminum derivatives

15

###### Catalyst solution 1:

Aluminum isopropoxide derived from diethylaluminum cyanide ( $\text{C}_2\text{H}_5)_2\text{AlCN}$ ) Into a 25-ml vial was added isopropanol (3 mmol) and toluene (9 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 1 ml of a 1 M solution of ( $\text{C}_2\text{H}_5)_2\text{AlCN}$  in toluene was added under nitrogen atmosphere.

20

###### Catalyst solution 2:

Aluminum isopropoxide derived from diethylaluminum cyanide ( $\text{C}_2\text{H}_5)_2\text{AlCN}$ )  
Similar procedure as for catalyst solution 1 followed by heating at 80°C for 1 h.

25

###### Catalyst solution 3:

Aluminum isopropoxide derived from triethylaluminum ( $(\text{C}_2\text{H}_5)_3\text{Al}$ ).  
Into a 25-ml vial was added isopropanol (4 mmol) and toluene (9 ml). The vial was

sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 1 ml of a 1 M solution of  $(C_2H_5)_3Al$  in toluene was added under nitrogen atmosphere.

The obtained catalyst solution was allowed to stand for 1 night at room temperature.

5

Catalyst solutions 1, 2 and 3 were used in standard racemisation protocol for (R)-2-amino-3-phenylpropionitrile:

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To the solution, catalyst solution was added by syringe and the reaction mixture was heated at temperature given in Table 5 below.

Table 5: Racemisation of (R)-2-amino-3-phenylpropionitrile using  $Al(O^iPr)_3$  derived from aluminum derivatives

| Exp. | Cat. sol | Catalyst (ml) | T (°C) | t (h) | e.e. (%) |
|------|----------|---------------|--------|-------|----------|
| 1    | 1        | 0.1           | 20     | 21    | 20       |
| 2    | 2        | 0.1           | 60     | 1     | 15       |
| 3    | 3        | 0.1           | 50     | 4     | 9        |
| 4    | 3        | 0.05          | 60     | 1     | 56       |
| 5    | 3        | 0.1           | 60     | 1     | 32       |

15

C) Aluminum isopropoxide prepared from other aluminum derivatives, catalytic racemisation in the presence of triethylamine

Catalyst solution:

20 Aluminum isopropoxide derived from trimethylaluminum  $((CH_3)_3Al)$ .

Into a 25-ml vial was added isopropanol (3 mmol) and toluene (9.5 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 0.5 ml of a 2 M solution of  $(CH_3)_3Al$  in toluene is added under nitrogen atmosphere.

25 The obtained catalyst solution was allowed to stand for 15 minutes at room temperature and for 15 minutes at 80°C respectively.

Stock solution triethylamine:

Into a 50-ml vial was added triethylamine (2,54 mmol) and toluene (25 ml). The vial



was sealed with a teflon-lined cap and stored at room temperature.

Catalyst solution was used in standard racemisation protocol for (R)-2-amino-3-phenylpropionitrile in presence of triethylamine:

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved  
 5 in toluene and stock solution of triethylamine was added. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To the solution, catalyst solution was added by syringe and the reaction mixture was heated at 60°C for 19 h.

10 Table 6: Racemisation of (R)-2-amino-3-phenylpropionitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$  derived from  $(\text{CH}_3)_3\text{Al}$  in the presence of triethylamine

| Exp. | Toluene (ml) | Catalyst solution (ml) | Stock triethylamine(ml) | e.e. (%) |
|------|--------------|------------------------|-------------------------|----------|
| 1    | 0.9          | 0.1                    | 0                       | 8        |
| 2    | 0.9          | 0                      | 0.1                     | > 99     |
| 3    | 0.8          | 0.1                    | 0.1                     | 27       |

D) Racemisation of (R)-2-amino-3-phenylpropionitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$  prepared from trimethylaluminum  $((\text{CH}_3)_3\text{Al})$  v.s. solid  $\text{Al}(\text{O}^i\text{Pr})_3$

15

Catalyst solution was derived from trimethylaluminum  $((\text{CH}_3)_3\text{Al})$ .

Into a 25-ml vial was added isopropanol (3 mmol) and toluene (9.5 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 0.5 ml of a 2 M solution of  $(\text{CH}_3)_3\text{Al}$  in toluene is added under nitrogen  
 20 atmosphere.

The obtained catalyst solution was allowed to stand for 15 minutes at room temperature and for 15 minutes at 80°C, respectively.

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) and solid  $\text{Al}(\text{O}^i\text{Pr})_3$  was dissolved in toluene. The vial was sealed with a teflon-lined cap and the  
 25 reaction mixture was degassed by nitrogen. To the solution, catalyst solution was added by syringe and the reaction mixture was heated at 60°C for 2 h. The results are given in Table 7.

Table 7: Racemisation of (R)-2-amino-3-phenylpropionitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$

| Exp. | Toluene (ml) | Catalyst solution (ml) | $\text{Al}(\text{O}^i\text{Pr})_3$ (mg) | e.e. (%) |
|------|--------------|------------------------|---|----------|
| 1    | 1            | 0                      | 3.4                                     | 72       |
| 2    | 0.9          | 0.1                    | 0                                       | 16       |

#### Example VI

- 5 Racemisation of (R)-2-amino-3-phenylpropionitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$  in the presence of trimethylsilyl cyanide (TMS-CN)

In a 5 ml vial, toluene (1 ml) was added to 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %),  $\text{Al}(\text{O}^i\text{Pr})_3$  (0.01 mmol) and TMS-CN. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. The temperature of the  
10 reaction mixture was increased to 80°C for 3 h. The results are given in Table 8.

Table 8: Racemisation of (R)-2-amino-3-phenylpropionitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$  in the presence of TMS-CN

| Exp. | TMS-CN (mmol) | e.e. (%) |
|------|---------------|----------|
| 1    | 0.1           | 23       |
| 2    | 0             | 56       |

15

#### Example VII

Racemisation of (R)-2-amino-3-phenylpropionitrile using diethylaluminum cyanide ( $\text{C}_2\text{H}_5)_2\text{AlCN}$ ) in combination with isopropanol

- 20 Catalyst solution:

Into a 50-ml vial was added toluene (10 ml). The vial was sealed with a teflon-lined cap and the toluene was degassed by nitrogen. To the toluene is added under nitrogen atmosphere respectively, 0.4 ml of a 1 M solution of  $(\text{C}_2\text{H}_5)_2\text{AlCN}$  in toluene and isopropanol (8.5 mmol).

- 25 The obtained catalyst solution was allowed to stand for 1 night at room temperature and used in catalytic racemisation of (R)-2-amino-3-phenylpropionitrile:

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 1 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture

was degassed by nitrogen. To the solution, catalyst solution was added by syringe and the temperature was increased to 80°C. The results are given in Table 9.

Table 9: Racemisation of (R)-2-amino-3-phenylpropionitrile using  
5 (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>AlCN/isopropanol (2eq.)

| Exp. | Catalyst solution (μl) | Catalyst (mol %) | t (h) | e.e. (%) |
|------|------------------------|------------------|-------|----------|
| 1    | 50                     | 2                | 6     | 37       |
| 2    | 250                    | 10               | 1     | 13       |

#### Example VIII

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using aluminum/amine catalyst

##### 10 Catalyst solution 1:

Into a 50-ml vial was added benzylamine (2 mmol) and toluene (9 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 1 ml of a 1 M solution of (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>AlCN in toluene was added under nitrogen atmosphere.

15

##### Catalyst solution 2, 3 and 4:

Into a 25-ml vial was added amine (3 mmol) and toluene (9 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 0.5

20 ml of a 2 M solution of (CH<sub>3</sub>)<sub>3</sub>Al in toluene was added under nitrogen atmosphere. Temperature was increased to 80°C for 1 night.

#### Catalyst solutions

| Catalyst solution | Amine                   |
|-------------------|-------------------------|
| 2                 | (S)-1-phenylethylamine  |
| 3                 | (RS)-1-phenylethylamine |
| 4                 | N-methylbenzylamine     |

25

Catalyst solutions were used in the catalytic racemisation of (R)-2-amino-3-phenylpropionitrile:

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To the solution, 0.1 ml catalyst solution was added by syringe and the temperature was increased to 60°C for 1 hour. The results are given in

5 Table 10.

Table 10: Racemisation of (R)-2-amino-3-phenylpropionitrile using aluminum/amine catalyst

| Catalyst solution | e.e. (%) |
|-------------------|----------|
| 1                 | 27       |
| 2                 | 13       |
| 3                 | 34       |
| 4                 | 10       |

10 Example IX

Racemisation of (R)-2-amino-3-phenylpropionitrile using diethylaluminum cyanide (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>AlCN in presence of isopropyl acetate

Catalyst solution:

15 Into a 50-ml vial was added toluene (9 ml). The vial was sealed with a teflon-lined cap and the solution was degassed by nitrogen. To the solution, 1 ml of a 1 M solution of (C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>AlCN in toluene was added under nitrogen atmosphere.

Catalyst solution was used in standard racemisation protocol for (R)-2-amino-3-phenylpropionitrile in presence of isopropyl acetate:

20 In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) and isopropyl acetate (0.11 mmol) were dissolved in 0,9 ml toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To the degassed solution, 0.1 ml catalyst solution was added by syringe and the reaction mixture was allowed to stand at ambient temperature for 1 h, after which 2-amino-3-phenylpropionitrile was racemised to 23 % e.e..

25

### Example X

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using lithium salts

LiBr was dried under vacuum at 300°C prior to use.

- 5 New LiPF<sub>6</sub> was used as such in glovebox.

#### Stock lithium salts

- 10 In a glovebox, lithium salt was added to a 25 ml-vial. The vial was sealed with a teflon-lined cap and taken out of the glovebox. Dry THF was added and the lithium salt was dissolved.

| Catalyst solution | Lithium salt (mmol) | THF (ml) |
|-------------------|---------------------|----------|
| LiBr              | 1,5                 | 15       |
| LiPF <sub>6</sub> | 1                   | 10       |

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using lithium salts

To a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was added.

- 15 The vial was sealed with a teflon-lined cap and degassed by nitrogen. To the (R)-2-amino-3-phenylpropionitrile-containing vial was added catalyst solution (1 ml) and the 2-amino-3-phenylpropionitrile was dissolved. The racemisation was performed under nitrogen at given temperature. The results are summarized in Table 11.

- 20 Table 11: Catalytic racemisation using lithium salts

| Exp. | Catalyst solution | T (°C) | t (h) | E.e. (%) |
|------|-------------------|--------|-------|----------|
| 1    | LiBr              | 80     | 24    | 34       |
| 2    | LiPF <sub>6</sub> | 60     | 2     | 40       |

### Example XI

#### Racemisation of (R)-2-amino-3-phenylpropionitrile nitrile using early transition metal catalysts

- 25

Racemisation using Zr(OBu<sup>t</sup>)<sub>4</sub>

Preparation catalyst solution, a general procedure:

Zr(OBu<sup>t</sup>)<sub>4</sub> was placed in a 5 ml vial under an atmosphere of dry nitrogen and toluene, freshly distilled from sodium metal, was introduced and the vial was sealed with a teflon-lined cap.

5

Table Zr(OBu<sup>t</sup>)<sub>4</sub> solutions in toluene

| Solution (M) | Zr(OBu <sup>t</sup> ) <sub>4</sub> (mmol) | Toluene (ml) |
|--------------|---|--------------|
| 0.1          | 1   | 10           |
| 1            | 2   | 2            |


Racemisation of (R)-2-amino-3-phenylpropionitrile

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml dry toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To this solution was added a toluene solution of Zr(OBu<sup>t</sup>)<sub>4</sub> (0.1 ml) by syringe and the reaction mixture was stirred for 24 h at 80°C. The results are given in Table 12.


15 Table 12: Catalytic racemisation of (R)-2-amino-3-phenylpropionitrile with Zr(OBu<sup>t</sup>)<sub>4</sub>

| Exp. | Catalyst solution (M) | E.e. (%) |
|------|-----------------------|----------|
| 1    | 0.1                   | 73       |
| 2    | 1                     | 0        |

Racemisation using different complexes



1



2

#### Catalyst solution

The complex (0.1 mmol) was placed in a 5 ml vial under an atmosphere of dry nitrogen and toluene (1 ml) freshly distilled from sodium metal was introduced. The vial was  
5 sealed with a teflon-lined cap and the mixture was stirred until the complex was completely dissolved.

#### Racemisation of (R)-2-amino-3-phenylpropionitrile.

10 In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml dry toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To this solution was added catalyst solution (0.1 ml) by syringe and the reaction mixture was stirred for 24 h at 50°C. The results are given in Table 13.

15

Table 13: Catalytic racemisation of (R)-2-amino-3-phenylpropionitrile

| Exp. | Complex | E.e. (%) |
|------|---------|----------|
| 1    | 1       | 6        |
| 2    | 2       | 75       |

#### Example XII

#### Racemisation of (R)-2-amino-3-phenylpropionitrile using aluminum alkoxides

20

Aluminum alkoxide catalysts prepared from  $\text{Me}_3\text{Al}$ .

Preparation catalyst solution, a general procedure:

Alcohol (3 mmol) was placed in a 25 ml-vial and toluene (9.5 ml) freshly distilled from sodium metal was introduced. The vial was sealed with a teflon-lined cap and  
25 degassed by nitrogen. To this solution was added a 2 M toluene solution of  $\text{Me}_3\text{Al}$  (0.5 ml, 1 mmol) and a stream of dry nitrogen removed the generated methane. After standing for 15 minutes at room temperature, the temperature is increased to 80°C for a period of time given in Table.

30 Racemisation of (R)-2-amino-3-phenylpropionitrile

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml dry toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To this solution was added catalyst solution (0.1 ml) by syringe and the temperature of the reaction mixture was increased to 60°C. Degrees of racemisation during a period of time are given in Table 14.

Table 14: Racemisation of (R)-2-amino-3-phenylpropionitrile catalysed by aluminum alkoxides

| Exp. | Preparation catalyst solution     |               | racemisation |          |
|------|-----------------------------------|---------------|--------------|----------|
|      | Alcohol                           | t (h) at 80°C | t (h)        | e.e. (%) |
| 1    | Isopropanol                       | 0.25          | 1            | 34       |
| 2    | (RS)-1-phenylethanol              | 0.25          | 1            | 77       |
| 3    | (S)-1-phenylethanol <sup>a)</sup> | 0.5           | 1            | 72       |
| 4    | 1-pentanol                        | 1             | 21           | 59       |
| 5    | 2-octanol                         | 0.5           | 1            | 44       |
| 6    | 2-cyano-2-propanol                | 0.25          | 1.25         | 74       |
| 7    | (RS)-1-cyclohexylethanol          | 1             | 1            | 55       |
| 8    | (RS)-1-(4-Chlorophenyl)ethanol    | 1             | 1            | 56       |
| 9    | (RS)-1-(4-Bromophenyl)ethanol     | 1             | 1            | 63       |

a) During catalyst preparation, e.e. of (S)-1-phenylethanol dropped to 54 %

10

Aluminum alkoxide catalyst prepared from <sup>i</sup>Bu<sub>3</sub>Al

Isopropanol (3 mmol) was placed in a 25 ml-vial and toluene (9 ml) freshly distilled from sodium metal was introduced. The vial was sealed with a teflon-lined cap and degassed by nitrogen. To this solution was added a 1 M toluene solution of <sup>i</sup>Bu<sub>3</sub>Al (1 ml, 1 mmol) and a stream of dry nitrogen removed the generated butane. After standing for 15 minutes at room temperature, the temperature is increased to 80°C for a period of time given in Table.

15

Racemisation of (R)-2-amino-3-phenylpropionitrile.

20

In a 5 ml vial, 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) was dissolved in 0.9 ml dry toluene. The vial was sealed with a teflon-lined cap and the reaction mixture was degassed by nitrogen. To this solution was added catalyst solution (0.1 ml)



by syringe and the temperature of the reaction mixture was increased to 60°C. Degree of racemisation is given in Table 15.

Table 15: Racemisation of (R)-2-amino-3-phenylpropionitrile catalysed by IPA/ $\text{Bu}_3\text{Al}$

| Exp. | Preparation catalyst solution |               | racemisation |          |
|------|-------------------------------|---------------|--------------|----------|
|      | Alcohol                       | t (h) at 80°C | t (h)        | e.e. (%) |
| 1    | Isopropanol                   | 0.25          | 1            | 61       |
| 2    | Isopropanol                   | 1.25          | 1            | 42       |

5

#### Example XIII

Racemisation of (R)-2-amino-3-phenylpropionitrile catalysed by bis(1,5-cyclooctadiene)rhodium (I) tetrafluoroborate

- 10 (R)-2-Amino-3-phenylpropionitrile (e.e. > 99 %) (146 mg, 0.1 mmol) and bis(1,5-cyclooctadiene)rhodium (I) tetrafluoroborate (4.5 mg, 0.011 mmol) were placed in a dry 5 ml vial. The vial was sealed with a teflon-lined cap, freshly distilled toluene (0.9 ml) was introduced and the reaction mixture was degassed by nitrogen. The temperature of the reaction mixture was increased to 60°C for 18 h, after which time (R)-2-Amino-3-phenylpropionitrile was racemised to 62 % e.e..
- 15

#### Example XIV

Racemisation of (R)-2-amino-3-phenylpropionitrile using metal isopropoxides

- 20 Catalyst solution  
Into a previously dried 25-ml vial was added metal isopropoxide (1 mmol) under an atmosphere of dry nitrogen. To the metal isopropoxide was added freshly from sodium distilled toluene (10 ml) and the vial was sealed with a teflon-lined cap.
- 25 Racemisation of (R)-2-amino-3-phenylpropionitrile  
A 5 ml vial was charged with 0.1 mmol (R)-2-amino-3-phenylpropionitrile (e.e. > 99 %) and dissolved in 0.9 ml dry toluene. The vial was sealed with a teflon-lined cap and the homogeneous solution was degassed by nitrogen. To this solution was added catalyst solution (0.1 ml) by syringe and the temperature of the reaction mixture was increased
- 30

to 60°C. Degrees of racemisation during a period of time using different metal catalysts are given in Table 16.

Table 16: Racemisation of (R)-2-amino-3-phenylpropionitrile catalysed by M(O<sup>i</sup>Pr)<sub>3</sub>

| Exp. | M(O <sup>i</sup> Pr) <sub>3</sub>  | racemisation |          |
|------|------------------------------------|--------------|----------|
|      |                                    | t (h)        | e.e. (%) |
| 1    | Ga(O <sup>i</sup> Pr) <sub>3</sub> | 1            | 85       |
| 2    | Sc(O <sup>i</sup> Pr) <sub>3</sub> | 5            | 92       |
|      |                                    | 21           | 77       |
| 3    | Y(O <sup>i</sup> Pr) <sub>3</sub>  | 5            | 46       |
|      |                                    | 21           | 23       |
| 4    | Yb(O <sup>i</sup> Pr) <sub>3</sub> | 24           | 54       |

5

#### Example XV

#### Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using Al(O<sup>i</sup>Pr)<sub>3</sub>

In two 5 ml vial, 0.1 mmol (R)-2-amino-2,3-dimethylbutyronitrile (e.e. 86 %) was weighed. To these vials 0.9 ml of toluene was added. The vials were sealed with teflon-lined caps and the solutions were degassed by nitrogen. To one of the degassed solution, 0.2 ml of a 0.1M Al(O<sup>i</sup>Pr)<sub>3</sub> catalyst solution in toluene was added by syringe. The temperature of the reaction mixture and blank reaction was increased to 60°C. After ½ hour a sample of the reaction mixture and blank was taken and analysed by HPLC. At this point the reaction mixture was racemic. The results are given in Table 17.

15

Table 17: Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using Al(O<sup>i</sup>Pr)<sub>3</sub>

| Exp. | Catalyst solution (ml) | t (h) | e.e. (%) |
|------|------------------------|-------|----------|
| 1    | 0                      | 0.5   | 86       |
| 2    | 0.2                    | 0.5   | <5       |

20

#### Example XVI

#### Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using AlEt<sub>3</sub>

In two 5 ml vial, 0.1 mmol (R)-2-amino-2,3-dimethylbutyronitrile (e.e. 86 %) was

weighed. To these vials 0.9 ml of toluene was added. The vials were sealed with teflon-lined caps and the reaction mixture and blank were degassed by nitrogen. To one of the degassed solutions, 0.2 ml of a 0.1M  $\text{AlEt}_3$  catalyst solution in toluene was added by syringe. The reaction mixture and blank reaction were allowed to stand at room temperature. After 5 minutes a sample of the reaction mixture and blank was taken and analysed by HPLC. At this point the reaction mixture was racemic. The results are given in Table 18.

Table 18: Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using  $\text{AlEt}_3$

| Exp. | Catalyst solution (ml) | t (min) | e.e. (%) |
|------|------------------------|---------|----------|
| 1    | 0                      | 5       | 86       |
| 2    | 0.2                    | 5       | <5       |

#### Example XVII

##### Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using $\text{Zr}(\text{tBuO})_4$

In two 5 ml vials, 0.1 mmol (R)-2-amino-2,3-dimethylbutyronitrile (e.e. 86 %) was weighed. To these vials 0.4 ml of toluene was added. The vials were sealed with teflon-lined caps and the reaction mixtures were degassed by nitrogen. To one of the degassed solutions, 0.6 ml of a 1M  $\text{Zr}(\text{tBuO})_4$  catalyst solution in toluene was added by syringe. The temperature of the reaction mixture and blank reaction was increased to 80 °C. After 1 day a sample of the reaction mixture and blank was taken and analysed by HPLC. At this point the reaction mixture was racemic. The results are given in Table 19.

Table 19: Racemisation of (R)-2-amino-2,3-dimethylbutyronitrile using  $\text{Zr}(\text{tBuO})_4$

| Exp. | Catalyst solution (ml) | t (h) | e.e. (%) |
|------|------------------------|-------|----------|
| 1    | 0                      | 24    | 86       |
| 2    | 0.6                    | 24    | <5       |

#### Example XVIII

##### Racemisation of (R)-2-amino-3-methylbutyronitrile using $\text{Al}(\text{O}^i\text{Pr})_3$

In two 5 ml vial, 0.1 mmol (R)-2-amino-3-methylbutyronitrile (e.e. 74 %) was weighed.

To both 0.9 ml toluene was added. The vials were sealed with a teflon-lined cap and the reaction mixtures were degassed by nitrogen. To one of the degassed solution, 0.1 ml of a 0.1M  $\text{Al}(\text{O}^i\text{Pr})_3$  catalyst solution in toluene was added by syringe. The temperature of the reaction mixtures was increased to 60°C. After 55 minutes a sample of the reaction mixture and the blank was taken and analysed by HPLC. At this point the reaction mixture was racemic. The results are given in Table 20.

Table 20: Racemisation of (R)-2-amino-3-methylbutyronitrile using  $\text{Al}(\text{O}^i\text{Pr})_3$

| Exp. | Catalyst solution (ml) | t (min) | e.e. (%) |
|------|------------------------|---------|----------|
| 1    | 0                      | 55      | 74       |
| 2    | 0.1                    | 55      | <5       |

10

#### Example XIX

#### Dynamic Kinetic Resolution of (R)-2-amino-3-phenylpropionitrile by racemisation with $\text{Al}(\text{O}^i\text{Pr})_3$ and enzymatic acylation catalyzed by Novozyme 435®

15 To a solution of 75 mg (0.51 mmol) (R)-2-amino-3-phenylpropionitrile and 104 mg (0.59 mmol) i-propyl phenylacetate in 4.2 g o-xylene was added 201 mg of Novozyme 435®. After stirring the reaction mixture for 20 minutes at ambient temperature, under reduced pressure (~100 mbar), 0.44 g (0.05 mmol) of a 0.1 M  $\text{Al}(\text{O}^i\text{Pr})_3$  solution in toluene was added. The reaction mixture was stirred at 70°C, under reduced pressure (20 ~100 mbar) for 43 hours. After ~20 hours another 185 mg Novozyme 435® and 0.42 g  $\text{Al}(\text{O}^i\text{Pr})_3$  solution in toluene was added. The reaction was analysed by HPLC. After 43 hours N-phenylacetyl-(S)-2-amino-3-phenylpropionitrile was formed 64% e.e.(S). The starting material was partially racemised.

#### 25 Example XX

#### Asymmetric transformation of 2-amino-3-phenylpropionitrile using $\text{Al}(\text{O}^i\text{Pr})_3$ as racemisation catalyst

The 1:1 diastereomeric salt of racemic 2-amino-3-phenylpropionitrile and N-(4-chlorobenzoyl)-L-glutamic acid was made by dissolving an 2.5 g (17 mmol) of 2-amino-3-phenylpropionitrile and 4.8 g (17 mmol) of N-(4-chlorobenzoyl)-L-glutamic acid in 68 ml of methanol. The mixture was stirred at room temperature till everything was

30

dissolved. The methanol was evaporated under reduced pressure to yield 7.2 g of the 1:1 diastereomeric salt (HPLC analysis: 2-amino-3-phenylpropionitrile < 5% e.e.).

In a Schlenk tube 0.860 g (2 mmol) of the 2-amino-3-phenylpropionitrile N-(4-chlorobenzoyl)-L-glutamic acid salt was dissolved in 4.8 ml toluene. This mixture was stirred and heated to 80°C overnight. Then, after cooling 0.2 ml of a 1M  $\text{Al}(\text{O}^i\text{Pr})_3$  solution in toluene was added. The reaction mixture was stirred and heated to 80°C for another 2 days. After cooling to room temperature the precipitate was filtered and washed with toluene. The residue (0.651 g) was analysed by HPLC. The results are given in Table 21. (Optical yield = yield x e.e.).

Table 21: Asymmetric transformation of 2-amino-3-phenylpropionitrile with  $\text{Al}(\text{O}^i\text{Pr})_3$

| Sample  | Yield (%) | e.e. (%) | Optical yield (%) |
|---------|-----------|----------|-------------------|
| Residue | 76        | 92 (R)   | 70                |

#### Example XXI

#### Asymmetric transformation of 2-amino-3-phenylpropionitrile in toluene using $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$ as racemisation catalyst

In 4.0 ml of methanol is dissolved 250 mg (1.7 mmol) of 2-amino-3-phenylpropionitrile, 480 mg (1.7 mmol) of N-(4-chlorobenzoyl)-L-glutamic acid and 100 mg of samarium iso-propoxide (  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  ). The methanol was removed under reduced pressure, the residue was suspended in 2 ml of toluene. The toluene was evaporated to remove the last traces of methanol and to the residue was added 4 ml of toluene. The reaction mixture was stirred for 21.5 h at 80°C. After cooling the diastereomeric salt and the  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  was filtered off. This residue was suspended in  $\text{CH}_2\text{Cl}_2$  and washed with saturated  $\text{NaHCO}_3$  solution. The  $\text{CH}_2\text{Cl}_2$  solution was dried and evaporated. The remaining (R)-2-amino-3-phenylpropionitrile (200 mg) was analysed by HPLC. The results are given in Table 22.

Table 22: Asymmetric transformation of 2-amino-3-phenylpropionitrile (Phe-CN) with  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  in toluene

| Sample | Yield (%) | e.e. (%) | Optical yield (%) |
|--------|-----------|----------|-------------------|
| Phe-CN | 73        | 90 (R)   | 66                |

Example XXII

Asymmetric transformation of 2-amino-3-phenylpropionitrile in heptane using  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  as racemisation catalyst

5

In 4.0 ml of methanol is dissolved 250 mg (1.7 mmol) of 2-amino-3-phenylpropionitrile, 480 mg (1.7 mmol) of N-(4-chlorobenzoyl)-L-glutamic acid and 100 mg of samarium iso-propoxide (  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  ). The methanol was removed under reduced pressure, the residue was suspended in 2 ml of heptane. The heptane was evaporated to remove the last traces of methanol and to the residue was added 4 ml of heptane. The reaction mixture was stirred for 21.5 h at 80°C. After cooling the diastereomeric salt and the  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  was filtered off. This residue was suspended in  $\text{CH}_2\text{Cl}_2$  and washed with saturated  $\text{NaHCO}_3$  solution. The  $\text{CH}_2\text{Cl}_2$  solution was dried and evaporated. The remaining (R)-2-amino-3-phenylpropionitrile (205 mg) was analysed by HPLC. The results are given in Table 23.

10  
15

Table 23: Asymmetric transformation of 2-amino-3-phenylpropionitrile (Phe-CN) with  $\text{Sm}_5\text{O}(\text{O}^i\text{Pr})_{13}$  in heptane

| Sample | Yield (%) | e.e. (%) | Optical yield (%) |
|--------|-----------|----------|-------------------|
| Phe-CN | 82        | 81 (R)   | 66                |

20

### CLAIMS

1. Process for the racemisation of an enantiomerically enriched  $\alpha$ -amino nitrile characterized in that the enantiomerically enriched  $\alpha$ -amino nitrile is contacted with a Lewis acid catalyst.  
5
2. Process according to claim 1, wherein an aprotic solvent is used.
3. Process according to claim 1 or 2, wherein the Lewis acid catalyst comprises a metal chosen from main group elements IA-IVA of the Periodic Table (CAS version), the transition metals and the lanthanides.
- 10 4. Process according to claim 3 wherein the metal is chosen from the group consisting of Al, Ti, Zr, or lanthanides.
5. Process according to any one of claims 1-4, wherein a catalyst with the general structure  $M_nX_pS_qL_r$  is used, wherein M represents the metal, X represents an anionic counterion or covalently bound anionic ligand for non zero valent metals, S represents a spectator ligand, L represents a neutral  
15 ligand, n represents an integer larger than or equal to 1 and p, q and r each independently represent an integer larger than or equal to 0, and in which n and p are chosen such that  $M_nX_p$  is neutral.
6. Process according to claim 5 wherein the catalyst is chosen from the group of  
20 aluminum alkoxides, aluminum alkyls, lanthanide alkoxydes and lanthanocenes.
7. Process according to any one of claims 1-6, wherein the racemisation is performed in combination with a resolution process.
8. Process according to claim 7, wherein the racemisation is performed in  
25 combination with an enzymatic resolution process.
9. Process according to claim 7, wherein the racemisation is performed in combination with a crystallization induced resolution.
10. Process according to any one of claims 7-9, wherein the resolution process is combined with racemisation in situ.
- 30 11. Process according to claim 10, wherein the racemisation is performed in situ in a crystallization induced asymmetric transformation process.

ABSTRACT

5        Process for the racemisation of an enantiomerically enriched  $\alpha$ -amino nitrile characterized in that the enantiomerically enriched  $\alpha$ -amino nitrile is contacted with a Lewis acid catalyst. Preferably an aprotic solvent is used.

      The Lewis acid catalyst preferably comprises a metal chosen from main group elements IA-IVA of the Periodic Table (CAS version), the transition metals and the lanthanides, in particular Al, Ti, Zr, or lanthanides.

10       The catalyst for example has the general structure  $M_nX_pS_qL_r$ , and preferably is chosen from the group of aluminum alkoxides, aluminum alkyls, lanthanide alkoxydes and lanthanocenes.

      The racemisation may be performed in combination with a resolution process, for instance in combination with an enzymatic or a crystallization induced resolution process, preferably in situ, for instance in situ in a crystallization induced  
15       asymmetric transformation process.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02412

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C253/30 C07C255/42 C07C255/24 C07C255/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No. |
|------------|--|-----------------------|
| X          | <p>LOPEZ-SERRANO P ET AL: "Enantioselective acylation of alpha-aminonitriles catalysed by Candida antarctica lipase. An unexpected turnover-related racemisation" TETRAHEDRON: ASYMMETRY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 12, no. 2, 19 February 2001 (2001-02-19), pages 219-228, XP004230902 ISSN: 0957-4166 page 220: second column, last paragraph; page 227: paragraphs 5.6.1-5.6.7</p> <p>---</p> <p>-/--</p> | 1-8,10                |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

10 February 2004

Date of mailing of the international search report

23/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fitz, W

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 2412

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| X          | <p>DATABASE WPI<br/>Derwent Publications Ltd., London, GB;<br/>AN 1993-383032<br/>XP002237430<br/>"IMPROVED RACEMISATION FOR<br/>2-AMINO:NITRILE(S) - COMPRISES CONTACTING<br/>OPTICALLY ACTIVE 2-AMINO:NITRILE(S) WITH<br/>POTASSIUM CYANIDE IN BENZENE-METHANOL<br/>MIXTURE"<br/>-&amp; JP 05 286919 A ((NIHA) NIKKO KYOSEKI<br/>KK), 2 November 1993 (1993-11-02)<br/>abstract</p> | 1-3,5                 |
| A          | <p>-----<br/>US 4 683 324 A (GASTROCK WILLIAM H ET AL)<br/>28 July 1987 (1987-07-28)<br/>cited in the application<br/>examples 1-3</p>  | 1                     |
| A          | <p>-----<br/>EP 1 050 529 A (AMERICAN CYANAMID CO)<br/>8 November 2000 (2000-11-08)<br/>cited in the application<br/>page 3, example 1<br/>-----</p>  | 1                     |

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 12412

| Patent document<br>cited in search report |   | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---|---------------------|----------------------------|---------------------|
| JP 5286919                                | A | 02-11-1993          | NONE                       |                     |
| US 4683324                                | A | 28-07-1987          | NONE                       |                     |
| EP 1050529                                | A | 08-11-2000          | AT 252549 T                | 15-11-2003          |
|   |   |                     | AU 3024500 A               | 09-11-2000          |
|   |   |                     | BR 0002074 A               | 02-01-2001          |
|   |   |                     | CA 2307392 A1              | 03-11-2000          |
|   |   |                     | CN 1272313 A               | 08-11-2000          |
|   |   |                     | DE 60006028 D1             | 27-11-2003          |
|   |   |                     | EP 1050529 A2              | 08-11-2000          |
|   |   |                     | HU 0001711 A2              | 28-05-2001          |
|   |   |                     | JP 2000319112 A            | 21-11-2000          |
|   |   |                     | NZ 504199 A                | 21-12-2001          |
|   |   |                     | PL 339972 A1               | 06-11-2000          |
|   |   |                     | SK 6012000 A3              | 18-01-2001          |
|   |   |                     | ZA 200002132 A             | 07-11-2000          |